

Manual

EPPiC

Kit for rapid, enzymatic purification of PCR products.

catalog #	size
1021-100	100 reactions
1021-500	500 reactions
1021-2500	2500 reactions

For research use only.

Guarantee

A&A Biotechnology provides guarantee on this product.

The company does not guarantee correct performance of this kit in the event of:

- not adhering to the supplied protocol
- use of not recommended equipment or materials
- use of other reagents than recommended or which are not a component of the product
- use of expired or improperly stored product or its components

Advantages

- Quick, lossless and effective PCR product purification.
- Better stability at room temperature compared to other SAP-based products.

Description

The EPPiC mixture contains two enzymes that effectively degrade dNTPs and primer left-overs from previous PCR mixtures while leaving the double stranded DNA PCR products untouched.

EPPiC enzymes are active at 37 °C in standard buffers used in PCR and are completely thermally inactivated by 15 min incubation at 70 °C.

The purified product can be used in downstream applications including nested, second round PCR, SNP analysis and cycle sequencing reactions.

Contents

	1021-100	1021-500	1021-2500	storage
EPPiC mixture	200 µl	1000 µl	5 x 1000 µl	-20 °C

Notes

Unlike the other enzymatic mixture used for PCR fragment clean up, EPPiC mixture does not remove 5'-phosphate groups from PCR products obtained with phosphorylated PCR primers. Therefore subsequent cloning of EPPiC purified phosphorylated PCR products does not require extra phosphorylation of 5'-ends.

Protocol

1. Briefly spin EPPiC mixture and place on ice.
2. To **10 μ l** of post-PCR reaction mixture add **2 μ l** of **EPPiC mixture**. Mix by pipetting.

Note. The reaction can be carried out with a different volume of post-PCR reaction mixture. However, a 5:1 ratio should be maintained (e.g. add 1 μ l of EPPiC mixture to 5 μ L of post-PCR reaction mixture).

3. Briefly spin the sample.
4. Incubate the sample in a thermal cycler:
 - 15 min at 37 °C
 - 15 min at 70 °C
5. Briefly spin the sample.
6. Store purified sample at -20 °C.



A&A BIOTECHNOLOGY
innovating life science

A&A Biotechnology, ul. Strzelca 40, 80-299 Gdańsk, Poland
phone +48 883 323 761, +48 600 776 268
info@aabiotech.com, www.aabiotech.com

